













JOINT TRAINING IN ATMOSPHERIC COMPOSITION


13 -17 OCTOBER 2025, BRUSSELS



How much will climate change this century?








observational constraints based on past simulated warming, as well as an updated assessment of equilibrium climate sensitivity. Very likely ranges are shown for the low and high GHG emissions scenarios (SSP1-2.6 and SSP3-7.0). **Right** - Global Reasons for Concern, comparing AR6 (thick embers) and AR5 (thin embers) assessments. Diagrams are shown for

Climate forcing, feedbacks, and sensitivity









Figure 7.10 | Global mean climate feedbacks estimated in abrupt4xCO2 simulations of 29 CMIP5 models (light blue) and 49 CMIP6 models (orange), compared with those assessed in this Report (red). Individual feedbacks for CMIP models are averaged across six radiative kernels as computed in Zelinka et al. (2020).



Net **cloud feedback** <u>positive</u> (*high confidence*).

Cloud feedbacks remain largest contributor to uncertainty of net feedback, but the uncertainty is reduced compared to AR5 (by ~50%). High confidence that radiative feedbacks are largest source of uncertainty in project global warming out to 2100 under increasing or stable emission scenarios.

Uncertainty is dominated by **aerosol ERF** in strong mitigation scenarios.

[chapter 7.5]

https://www.ipcc.ch/report/ar6/wg1/

Figure TS.14 | Schematic representation of changes in the top-of-atmosphere (TOA) radiation budget following a perturbation. The intent of this figure is to

Climate and cloud feedbacks (AR6 7.4.2.4)







Figure 7.9 | Schematic cross section of diverse cloud responses to surface warming from the tropics to polar regions. Thick solid and dashed curves indicate the tropopause and the subtropical inversion layer in the current climate, respectively. Thin grey text and arrows represent robust responses in the thermodynamic structure to greenhouse warming, of relevance to cloud changes. Text and arrows in red, orange and green show the major cloud responses assessed with high, medium and low confidence, respectively, and the sign of their feedbacks to the surface warming is indicated in the parenthesis. Major advances since AR5 are listed in the box. Figure adapted from Boucher et al. (2013).

Cloud feedback A *climate feedback* involving changes in any of the properties of clouds as a response to a change in the local or global surface temperature. Understanding cloud feedbacks and determining their magnitude and sign requires an understanding of how a change in *climate* may affect the spectrum of cloud types, the cloud fraction and height, the radiative properties of clouds, and finally the Earth's radiation budget.

Box SPM.1.3: The CMIP6 models considered in this Report have a wider range of climate sensitivity than in CMIP5 models and the AR6 assessed *very likely* range, which is based on multiple lines of evidence. These CMIP6 models also show a higher average climate sensitivity than CMIP5 and the AR6 assessed best estimate. The higher CMIP6 climate sensitivity values compared to CMIP5 can be traced to an amplifying cloud feedback that is larger in CMIP6 by about 20%. {Box 7.1, 7.3, 7.4, 7.5, TS.3.2}



FAQ 7.2, Figure 1 | Interactions between clouds and the climate, today and in a warmer future. Global warming is expected to alter the altitude (left) and the amount (centre) of clouds, which will amplify warming. On the other hand, cloud composition will change (right), offsetting some of the warming. Overall, clouds are expected to amplify future warming.

FAQ 7.2 | What Is the Role of Clouds in a Warming Climate?

One of the biggest challenges in climate science has been to predict how clouds will change in a warming world and whether those changes will amplify or partially offset the warming caused by increasing concentrations of greenhouse gases and other human activities. Scientists have made significant progress over the past decade and are now more confident that changes in clouds will amplify, rather than offset, global warming in the future.

Clouds cover roughly two-thirds of the Earth's surface. They consist of small droplets and/or ice crystals, which form when water vapour condenses or deposits around tiny particles called *aerosols* (such as salt, dust, or smoke). Clouds play a critical role in the Earth's *energy budget* at the top of our atmosphere and therefore influence Earth's surface temperature (see FAQ 7.1). The interactions between clouds and the climate are complex and varied. Clouds at low altitudes tend to reflect incoming solar energy back to space, creating a cooling effect by preventing this energy from reaching and warming the Earth. On the other hand, higher clouds tend to trap (i.e., absorb and then emit at a lower temperature) some of the energy leaving the Earth, leading to a warming effect. On average, clouds reflect back more incoming energy than the amount of outgoing energy trap, resulting in an overall net cooling effect on the present climate. Human activities since the pre-industrial era have altered this climate effect of clouds in two different ways: by changing the abundance of the aerosol particles in the atmosphere and by warming the Earth's surface, primarily as a result of increases in greenhouse gas emissions.

earthcare earthcare

CLOUDS AND CLIMATE



PRESENT CLIMATE

High thin clouds trap outgoing heat more than they reflect incoming solar energy, warming the atmosphere

FUTURE CLIMATE

More high thin clouds mean more outgoing heat is trapped

Fewer low thick clouds mean less solar energy is reflected back to space.

Cloud distribution is expected to change in the future and amplify climate warming



CLIMATE


Clouds affect and are affected by climate


Low thick clouds reflect **incoming solar energy** more than they trap outgoing heat, cooling the atmosphere

Overall, clouds currently cool the climate system

ERF and aerosol-cloud-radiation interactions









Effective radiative forcing of aerosols and clouds







7.3.3.4. In AR5, the overall assessment of total aerosol ERF (ari+aci) used the median of all ESM estimates (...) but placed more confidence in a subset of models that were deemed more complete in their representation of aerosol-cloud interactions. These models, which include aerosol effects on mixed-phase, ice, and/or convective clouds, (...)

Likewise, studies that constrained models with satellite observations (...) were given extra weight. (...)

Finally, based on higher resolution models, doubt was raised regarding the ability of ESMs to represent the cloud-adjustment component of ERFaci with fidelity.

[In AR6], the best estimate and range is revised relative to AR5, partly based on updates to the above lines of argument. (...) As in AR5, there is still reason to question the ability of ESMs to simulate adjustments in LWP and cloud cover in response to aerosol perturbation, (...)


There is thus strong evidence for a substantive negative total aerosol ERF, which is supported by the broad agreement between observation-based and model-based lines of evidence for both ERFari and ERFaci that has emerged since AR5.

However, considerable <u>uncertainty</u> remains, particularly with regards to the adjustment contribution to ERFaci, as well as <u>missing processes</u> in <u>current ESMs</u>, notably aerosol effects on <u>mixed-phase</u>, ice, and convective clouds. This leads to a medium confidence in the estimate of ERFari+aci and a slight narrowing of the uncertainty range.

There is high confidence that ERFaci contributes most (75-80%) to the total aerosol effect.

Next-gen climate models with interactive aerosols (ICON-HAM-lite)


Weiss et al., 2025, GMD (5 km model grid)



AEROSOLS AND CLIMATE

esa

Aerosols – tiny particles such as dust suspended in the air – affect the climate, either directly necting solar energy back to space and by trapping heat in the atmosphere, or indirectly by influencing how clouds form







EarthCARE: an Earth Explorer satellite mission for cloud, aerosol, and Earth radiation budget research


09/10/2025

Alex Hoffmann on behalf of the joint ESA and JAXA EarthCARE Team

Joint Training in Atmospheric Composition Online Webinar


ESA's Earth Observation fleet





Earth Explorers:







From early ideas to implementation...



The Earth Explorer Earth Radiation Mission A Tobias, R Meynart, P Ingmann, W Leibrandt ESA/ESTEC - Noordwijk - The Netherlands

The Earth Explorer Earth Radiation Mission proposed by the scientific commanity for the ESA Earth Explorer line of missions will provide for the first time a multi-year set of cloud profiling and aerously observations to progress in understanding the interaction of congress of water between the Earth's safface and the cycle of the attemporter. A backconter lister and a cloud profiling part will be used separately, in complement and space gost provide these profiles that will be contained across task by a visible-started colon language and a row-destined brookbast radiomate, from a satellite at eval the Orthodol across task by a visible-started colon language and a row-destined brookbast radiomate, from a satellite at eval the Orthodol across task by a visible-started colon language and a row-destined brookbast radiomate, from a satellite at eval the Orthodol across task by a visible-started colon language and a row-destined brookbast radiomate, from a satellite at eval the Orthodol across task by a row of the row of the orthodol across task by a row of the row of the orthodol across task by a row of the is unique for the combination and capability of its sensors and for its utilisation

The Earth Radiation Mission (ERM) is one of the four candidates for implementation in the research oriented Earth Explorer line of missions within ESA's new T-living Flaset* programme* which also includes the Earth Watch line of service-oriented missions. The ERM is competing with three other candidates, the Gravity-field and steady-state Ocean Circulation Explore, the Land-Surface Processes and Interactions Mission and the Autonosheric Dynamics Mission in a process that should cultimiste in missions. The EBM is competing with three other candidates, the Gravity-field and study-state Dorsan Circuitation Explorer, the Land-Surface Processom and Interactions Missions and the Attrosphority Dynamics Mission in a process that should colimitate in Land-Surface Processom and Interactions Missions and the Attrosphority Dynamics Mission in process that should colimitate in understaken radials and development starvities with the European and Canadians Surface Community and industry. This page summarises the presert results. Each of these activities were presented recently at 8 canadian of Radiation Sudice organized by ESA's Earth Observation Programme. It would be impressed only in the references all scientific and technical reports, each of the Canadian Sudice Sudi

2. MISSION JUSTIFICATION AND ORJECTIVES

The need for the ERM [2],[3] can be intuitively explained with table 2.1, by Chahine. The effects of uncertainties in our capabilities to estimate clouds and hamidity in the atmosphere are much larger than the effect of the CO₂ which is however most commonly known in the media.

Parameter	Current uncertainty	Effect on downwelling infrared flux Wm ⁻¹	
		Sub-artic summer	Tropics
Specific humidity	20%	11	15
Atm. temperature	2 K	5	8
Cloud base height	100 hPa	11	7
Fract.cloud cover	0.1	6	3
Doubling of CO ₂ with 50% cloud cover, cloud base at 400 hPa		0.8	0.2
Doubling of CO, with an about source on find healt		1.2	0.2


ver, no feed-back 1.2
Table 2.1. The need for the ERM (table after Chahine)

ctive of the ERM, schematically shown in figure 2.1, is therefore to progress in understanding the transport of energy between the Earth's surface and the top of the atmosphere.

and the Atmosphere III • Barcelona, Spain • September 1998 SPIE Vol. 3495 • 0277-786X/98/\$10.00







https://doi.org/10.5194/amt-16-3581-2023

O Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.

Measurement EGU Techniques

The EarthCARE mission - science and system overview

Tobias Wehr1.7, Takuji Kubota2, Georgios Tzeremes1, Kotska Wallace1, Hirotaka Nakatsuka2, Yuichi Ohno Rob Koopman¹, Stephanie Rusli¹, Maki Kikuchi², Michael Eisinger³, Toshiyuki Tanaka², Masatoshi Taga², Patrick Deghaye¹, Eichi Tomita², and Dirk Bernaerts¹

sun-yean пракс «детох, ISA»—ISTES, Keplerlant I, 2201 AZ Noordoi); the Netherlands "Japan Accouptes Epipelarian Agency (JAAN), 305-8505 (Amee-1-1, Senger, Tsukuba, Baraki, Japan "Вигореая Брасс Адено, ISA»—ISCAN, Fermi Averase, Didco, (XXI 10FD, UK "Asiacaal Institute of Information and Commissications Technology (NICT), 184-879-4-2-1 Nakai-Katamachi, Koganet, Tokyo, Japan "Accessed, 1 Peterusy 2023" ¹ Furonean Space Agency, ESA - ESTEC, Kenlerlaan 1, 2201 AZ Noordwijk, the Netherlands

Correspondence: Kotska Wallace (kotska.wallace@esa.int) and Takuji Kubota (kubota.takuji@jaxa.jp)

Received: 22 December 2022 - Discussion started: 21 February 2023 Revised: 9 May 2023 - Accepted: 9 May 2023 - Published: 4 August 2023

Abstract, The Earth Cloud, Aerosol and Radiation Explorer Abstract. The Earth Cloud, Aerosol and Radiation Explorer (EarthCARE) is a staellite mission implemented by the Eu-ropean Space Agency (ESA), in cooperation with the Japan Aerospace Exploration Agency (JAXA), to measure global profiles of aerosols, clouds and precipitation properties to-gether with radiative fluxes and derived heating rates. The imultaneous measurements of the vertical structure and horzontal distribution of cloud and aerosol fields, together with outgoing radiation, will be used in particular to evaluate their representation in weather foresteading and climate models and representation in weather foresteading of cloud and aerosol radia-tive impact and feedback mechanisms. To achieve twice, the goal is she activitied secure with footprint size of twice, the goal is she activitied secure with footprint size of 10 km × 10 km is measured with sufficiently of the twice with the state of th

To optimise the performance of the two active instrument the platform will fly at a relatively low altitude of 393 km. with an equatorial revisit time of 25 d. The scientific payloa consists of four instruments: an atmospheric lidar, a cloud-profiling radar with Doppler capability, a multi-spectral im-ager and a broadband radiometer. Co-located measurements from these instruments are processed in the ground segment, which produces and distributes a wide range of science data products. As well as the Level 1 (L1) product of each inhave been developed, in both Europe and Japan, benefiting

ment with a detailed description of the four science instru ments, demonstrating how the observational requirement will be met. Furthermore, the elements of the space seemen and ground segment that are relevant for science data user

The Earth Cloud, Aerosol and Radiation Explorer (Earth The Earth Cloud, Aerosol and Radiation Explorer (Earth-CARE) will provide global profiles of clouds, aerosols and precipitation along with co-located radiative flux measure-ments. Atmospheric microphysical properties and associ-ated radiative fluxes will be used to evaluate the representa-tion of aerosols, clouds and precipitation in weather forecast

The EarthCARE satellite includes four scientific instruments: an atmospheric lidar, a Doppler cloud radar, a multi-spectral imager and a broadband radiometer. The instruments have been characterised, calibrated, tested and integrated on

2001 2004 1998

> EarthCARE Candidate (Ph A) Report for Mission Selection

2023

Earth Radiation Mission Candidate https://doi.org/10.1117/12.332702

> EarthCARE Candidate (Ph 0) Report for Assessment

Science and system overview publication Illingworth et al, BAMS

Mission description publication https://doi.org/10.1175/BAMS-D-12-00227.1

https://doi.org/10.5194/amt-16-3581-2023









































2015





Wehr et al, AMT









EarthCARE: Scientific objectives





Systematic provision of vertical profiles of clouds and aerosols, collocated with measurements of solar affow do aerosols and radiation clouds heat or cool the Direct ver Heating of impact of clouds and aerosols on atmospheric heating rates and radiative fluxes.


See EarthCARE Mission Requirements Document (MRD) for more details on mission objectives and requirements.





EarthCARE: instrument payload







EarthCARE: key measurement specs






Cloud Profiling Radar CPR


High power 94GHz cloud radar with Doppler

- 2.5 m antenna
- Resolution 800m (hor) x 500m (ver)
- Range -1 km to 20 km
- Doppler velocity range ±10ms⁻¹
- Doppler velocity accuracy 1ms⁻¹ @10km integration, -19dBZ

L1: radar reflectivity

& vertical Doppler velocity





Multi Spectral Imager MSI

- 150km swath (-35 km to +115 km)
- 500m Ground Sampling Distance
- VIS, Near IR, SWIR Camera (VNS)
- Thermal IR Camera (TIR) L1: radiances (VNS). brightness temperature (TIR)



Broad Band Radiometer BBR

- TOA fluxes with accuracy better than 10Wm⁻²
- 3 telescopes, Total Wave and Short Wave radiances at same location, different angles
- SW 0.25 to 4 μm, LW 4 μm to >50 μm
- Angular sampling 0, +/- 55°
- Spatial resolution 10kmx10km
- Spatial sampling 1 km
- L**1**: Solar and Thermal TOA rad.

Atmospheric LIDAR ATLID

High spectral resolution, **UV LIDAR**

- 355 nm source
- Vertical profiles 0-40km, 100-500m resolution
- Horizontal sampling 280m
- Rayleigh co-pol
- Mie co-pol
- Mie cross-pol

L1: attenuated backscatter profiles







CPR instrument



- 94 GHz cloud radar with Doppler capability
 - 2.5 m transmit/receive antenna (800 m footprint)
- Transmitter
 - Emits 3.3 μs pulses, peak power 1500 W
- Receiver
 - Vertical resolution 500 m oversampled at 100 m
 - 500 m horizontal integration
 - 3 Pulse Repetition Frequencies, dependant on latitude, 6100 to 7500 Hz
 - Minimum sensitivity better than -35 dBZ for 10km integration and uniform clouds
 - Doppler velocity measurement range $\pm 10~{
 m ms}^{-1}$
 - Doppler velocity measurement accuracy 1 ms⁻¹
 @10km integration, -19 dBZ




- → Profiles of radar reflectivity, vertical velocities of cloud and precipitation particles.
- → Radar signal allows penetration much deeper into clouds, compared to lidar.
- → Synergistic use with the atmospheric lidar and multispectral imager used to classify atmospheric targets and derive atmospheric profiles, important for making radiative transfer calculations.




ATLID instrument




- Bi-static, 355 nm UV lidar
 - Actively controlled co-alignment loop
- Transmitter
 - Power laser head emits **37 mJ** pulses of 26 ns at a repetition rate of 51 Hz
 - Linear polarised output
 - 15 m ground footprint points 3° backwards to minimise specular reflection
- 3-channel receiver
 - Cross-polar channel
 - Co-polar, spectrally broadened molecular backscatter ("Rayleigh") channel
 - Co-polar, particulate backscatter channel ("Mie")





Molecular backscatter contribution






- → Aerosol properties including altitude, thickness, detailed optical properties and aerosol type.
- → High-spectral resolution receiver and depolarisation channel.



MSI instrument



- 7 channel imager
- Visible, Near-IR and Shortwave IR camera
 - 4 separate optic trains
 - Pushbroom, linear detectors
- Thermal IR camera
 - Three channels share defined by dichroics
 - Common 2-D detector uses Time Delay Integration to increase SNR




- → Provision of scene context information.
- → Derivation of "MSI-only" cloud and aerosol data products.
- → Nadir pixel used in retrieval of synergistic products with the ATLID and CPR.
- → Across-track data (selected channels) used for creation of 3D scenes, for 1D and 3D radiative transfer calculations to calculate radiative properties.
- → Used to improve the unfiltering of BBR measured radiances.



BBR instrument



- Broadband Radiometer
 - 3 views: forewards, aft and nadir
 - 2 channels per view: short-wave and total-wave, from which is derived a long-wave
 - Pushbroom onto microbolometer arrays
 - Unfiltering for the spectral response of the scene is improved using data from the MSI





→ Unfiltered, top-of-atmosphere radiances, short-wave and longwave fluxes





Data processing chain



CPR Level 1b (JAXA)

- Radar reflectivity
- Doppler velocity profiles

ATLID Level 1b (ESA)

Attenuated backscatter in

- Rayleigh channel
- Co-polar Mie channel
- Cross-polar Mie channel

MSI Level 1b/c (ESA)

TOA radiances for four solar channels, TOA brightness temperatures for three thermal channels

BBR Level 1b (ESA)

Filtered TOA short-wave and total-wave radiances

CPR Level 2a

Radar echo product, feature mask, cloud type, liquid and ice cloud properties, vertical motion, rain and snow estimates, ...

ATLID Level 2a

Feature mask and target classification, extinction, backscatter & depol. profiles, aerosol properties, ice cloud properties, ...

MSI Level 2a

Cloud mask, cloud microphysical parameters, cloud top height, aerosol parameters, ...

BBR Level 2b

Unfiltered top-of-atmosphere radiances, short-wave and long-wave fluxes Products are enhanced by using MSI



Target classification, cloud &



aerosol prof. at x-sectn, ...



3D Scenes Construction


Expand syn. retrievals acrosstrack using MSI; ≈40km wide

microphsyical observations + theory

Assessment

Comparison of calculated fluxes and radiances to BBR observations

direct observations



Operational software developed directly by science teams

Radiative Transfer Products calculated radiances, fluxes, heating rate profiles






EarthCARE data production model (ESA)










AMT special issue: *EarthCARE Level 2 algorithms and data products* https://amt.copernicus.org/articles/special_issue1156.html



Atmos. Meas. Tech., 17, 3081–3083, 2024 https://doi.org/10.5194/amt-17-3081-2024 © Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Preface to the special issue "EarthCARE Level 2 algorithms and data products": Editorial in memory of Tobias Wehr

Robin J. Hogan^{1,2}, Anthony J. Illingworth², Pavlos Kollias³, Hajime Okamoto⁴, and Ulla Wandinger⁵

- ¹European Centre for Medium-Range Weather Forecasts, Reading, UK
- ²Department of Meteorology, University of Reading, Reading, UK
- ³School of Marine and Atmospheric Sciences, Stony Brook University, New York, USA
- ⁴Research Institute for Applied Mechanics, Kyushu University, Fukuoka, Japan
- ⁵Leibniz Institute for Tropospheric Research, Leipzig, Germany

Correspondence: Robin J. Hogan (r.j.hogan@ecmwf.int)

Received: 30 April 2024 - Published: 22 May 2024








EarthCARE data products











- Mutual dependency at Level 1 requiring for strong coordination between Agencies
- Separate Level 2 chains provide opportunity for collaboration and comparison
- Each agency coordinates the validation of its own products



Data release schedule



March 2025

Level 1b Single Sensor Products


Level 2a Single Sensor Products

Level 2b Two Sensor Products

Level 2b Synergistic Products



https://earth.esa.int/eogateway/missions/earthcare/data



https://www.esa.int/Applications/Observing_the_Earth/FutureEO/EarthCARE/EarthCARE_goes_live_with_data_now_available_to_all




Synergy images,






Atmospheric heating

from EarthCARE in synergy



https://www.esa.int/Applications/Observing the Earth/FutureEO/EarthCARE/EarthCA

RE synergy refeals power of clouds and aerosols








EarthCARE validation sources







Airborne campaigns
Under-flights for product
validation:
Rapidly collect numerous,
precise collocations.




Networks

Continuous surfacebased data acquisition Slower collection of collocations but broader coverage of geophysical and meteorological conditions.













EarthCARE data with aircraft tracks



Atlantic Ocean






https://www.esa.int/Applications/Observing_the_Earth/FutureEO/EarthCARE/Taking_to_the_skies_for_EarthCARE

Cloud structure measured by Cloud Profiling Radar (CPR) 💸

ATLID: atmospheric lidar fundamentals







range $P_L(R) = P_o \eta_T \eta_R G(R) \left(\frac{A}{R^2}\right) \left(\frac{ct}{2}\right) \beta(R) \exp\left[-2 \int_0^R \alpha(R') dR'\right]$

Laser power per pulse

system variables

atmospheric variables

molecules

aerosols

B: volume backscatter coefficient

α: extinction coefficient: absorption + scattering by molecules + aerosols



High spectral resolution, **UV LIDAR**

- 355 nm source
- Vertical profiles 0-40km, 100-500m resolution
- Horizontal sampling 280m
- Rayleigh co-pol
- Mie co-pol
- Mie cross-pol

L1: attenuated backscatter profiles










ATLID: High Spectral Resolution Lidar







Compared to e.g. CALIPSO where total (Mie+Ray) signals are not separated:

- Direct measurement of aerosol extinction
- Measurement of extinction-to-backscatter ratio
- Improved daylight performance

Non-HSRL lidar

One measurement but two unknowns

$$P(z) = \frac{1}{z^2} \left(\beta_{Part}(z) + \beta_{Mol}(z) \right) e^{-2\int_0^z \alpha(z')dz'}$$

Must assume $S = \frac{\alpha_{Part}}{1}$ in order to

recover backscatter and extinction.

HSRL lidar

Two measurement and two unknowns

$$P_{Part}(z) = \frac{1}{z^2} \beta_{Part}(z) e^{-2\int_0^z \alpha(z')dz'}$$

$$P_{Mol}(z) = \frac{1}{z^2} \beta_{Mol}(z) e^{-2\int_0^z \alpha(z')dz'}$$
$$= \sigma_{Ray} \frac{1}{z^2} N(z) e^{-2\int_0^z \alpha(z')dz'}$$


→ Can determine S!



ATLID: HSRL and polarization receiver








ATLID has three channels:

- A perpendicular channel
- 2. A parallel (mainly-) particulate scattering channel
- 3. A parallel (mainly-) molecular scattering channel




After spectral/polarization cross-talk correction and calibration

- 1. Perpendicular particulate channel (cross-polar Mie)
- 2. Parallel particulate scattering channel (co-polar Mie)
- 3. Molecular scattering channel (Rayleigh)




ATLID: Level-1 data





- **High spectral resolution lidar (HSRL)** separately measures backscatter of particles (clouds & aerosols) and air (Rayleigh scattering)
- Can unambiguously calculate the important profile of extinction coefficient from how much the air backscatter is attenuated

Aerosols modelled with NASA's Goddard Earth Observing System






ATLID: aerosol type classification








Aerosol classification from measurements of lidar ratio and particle linear depolarization ratio at 355 nm.

Ground-based observations were performed with the Raman-polarization lidars (POLIS) (University of Munich, dots) and Polly XT (Leibniz Institute for Tropospheric Research, open squares) at Cape Verde (dust, marine, dust and smoke, dusty mixtures; dots; Groß et al. 2011); Leipzig, Germany (pollution, aged boreal biomassburning aerosol, dusty mixtures; open squares); Munich, Germany (volcanic ash; dots; Groß et al. 2012); in the Amazon basin (smoke; open squares; Baars et al. 2012); and over the North Atlantic (dust, dust and smoke; open squares; Kanitz et al. 2013). Source: Figure 8, Illingworth et al. 2015



ATLID: A-PRO cloud and aerosol processor





A-TC (target classification algorithm), description in: Donovan et al., 2024, Wandinger et al., 2023 and Irbah et al., 2023



ATLID: smoke from wildfires into UTLS





https://earth.esa.int/eogateway/success-story/earthcare-sees-canadian-wildfire-smoke-spreading-over-the-northern-hemisphere


2025 was second only to 2023 in the intensity of the Canadian wildfires, and just ahead of 2024.


Injection of dense smoke plumes into the upper atmosphere.

EarthCARE can tracked smoke layers over vast distances.

What's the impact on long-term climate, ozone,

cloud formation and radiative forcing?







Towards scientific exploitation






https://earth.esa.int/eogateway/missions/earthcare

EarthCARE Product Data Handbook: https://earthcarehandbook.earth.esa.int/

https://www.earthcarescience.net/home



New measurements

What can we learn from the Doppler radar and the high spectral resolution lidar?




Radiative closure

Radiation calculations check that retrieved clouds are consistent with broadband radiation measurements.



Model evaluation

How can we use the data to evaluate and improve weather and climate models?



Data assimilation

How will the data be ingested into weather forecast and airquality models?






Radiative closure




https://www.earthcarescience.net/science/radiative-closure

https://www.ecmwf.int/en/about/media-centre/news/2025/earthcare-instruments-working-together-evaluate-atmospheric-models



- EarthCARE continually compares computed fluxes from ACM-RT to measurements by the broadband radiometer (processed by BMA-FLX)
- If they agree, it gives us confidence in the retrieved cloud properties; if they disagree then we have something to improve in our algorithms or something to learn about the atmosphere!





Model eval: CAMS aerosols extinction

ATLID A-EBD

Peter Hill, Robin Hogan, David Donovan

ATL-EBD particle extinction coefficient 355nm low resolution; aerosol only





https://www.earthcarescience.net/science/model-evaluation


CAMS: ECMWF's air quality forecast including 15 prognostic aerosol variables


Outputs 355-nm extinction coefficient

Have never before been able to evaluate aerosol profiles globally with such resolution & accuracy, nor above clouds!














Model eval: CAMS aerosols in India

Peter Hill, Robin Hogan, David Donovan











Improving ice/snow fall speed in ECMWF model



CPR mean fall speed

EarthCARE

---- Mean 0°C level

3.0


2.5

1.5 © Ice & snow for climate and we have been some for the state of the state of


Rebecca Murray-Watson, Mark Fielding, Richard Forbes

Ice & snow fall speeds affect model climate and forecast skill: this is first time we have been able to evaluate it globally!


ECMWF model is very simple! ice falls at 0.13 m/s, snow at 1 m/s




EarthCARE



Modifications: use a mass-weighted fall speed; add rimed snow, temperature-dependent size & air-density effect








Data assimilation and NWP impact




https://www.ecmwf.int/en/newsletter/183/earth-system-science/earthcare-data-begin-make-impact



Assimilation of CPR reflectivity and ATLID backscatter into ECMWF model improves skill: *will go operational in Cycle 50r1 (autumn 2025)* ECMWF is working on assimilation of ATLID into aerosol forecasts

Impact on 5-day forecast skill






Gravity waves in Antarctic cloud




https://earth.esa.int/eogateway/success-story/earthcare-reveals-how-atmospheric-ripples-boost-cloud-formation-over-antarctica

Gravity waves are common in the atmosphere, particularly in the vicinity of orography, and can lead to cloud formation.


In this example over Antarctica, gravity waves were detected by the CPR with a wavelength of 18 km and an amplitude up to 2 m/s.






Polar Stratospheric Clouds







- Polar Stratospheric Clouds (PSCs) over the Arctic (and Antarctica)
- Form inside the polar vortex, at heights of ~20-30 km
 - PSC Type I: supercooled nitric acid (and sulfuric acid) and water (T < -78°C)
- Contribute to polar ozone depletion (see <u>Tritscher et al., 2021</u>)
- PSC Type II: ice crystals (nacreous clouds) ($T \sim < -83$ to -85 °C)
- Mountain leewaves (gravity waves) can form lenticular PSCs



A direct look into the eye of a hurricane





https://www.earthcare science.net/gallery/tro pical-cyclones See also:


https://earth.esa.int/eo gateway/successstory/earthcare-staresinto-the-eye-oftropical-cyclone-vince













M-REFF Cloud Effective Radius



M-CTH Cloud Top Height [km]















Observing volcanic plumes



https://www.earthcarescience.net/gallery/volcanoes







In a nutshell:



- EarthCARE explores the scientific basis of the interactions between cloud, aerosol, and radiation, and will
 improve satellite algorithms, numerical weather predictions, and climate model projections, with new features
 such as synergetic observations by four sensors
- EarthCARE hosts two unique remote sensing instruments, CPR (Doppler Radar) and ATLID (High Spectral Resolution Lidar)
- The EarthCARE satellite was successfully launched on 28 May 2024
 - Level 1 products were released to the public on 14th January 2025
 - Level 2 single-sensor products and 2-sensor synergy products were released on 17th March 2025
 - 3-sensor and 4-sensor synergy products are scheduled to be released in December 2025
- Validation results show very promising early product quality and offering significant scientific opportunities.










BACK-UP SLIDES




Evolving ERF(ari+aci) uncertainties












1998 > 2001

2004

2015

2023






Figure 2.1: Global annual mean radiative forcing at the top of atmosphere corresponding to changes from 1750 to 2000 following IPCC (2001) together with some more recent assessments of aerosol direct and indirect effects (modelling and satellite studies). The IPCC values show smaller direct forcings than recently derived by Bellouin et al. (up to -1.9 Wm², see text). Note that although direct radiative forcing is fairly well known, indirect forcing of aerosols is extremely uncertain. (A: Takemura et al., 2002, B: Kaufman et al., 2003, C: Takemura, 2003a, D: Takemura, 2003b, E: Nakajima et al., 2001, F: Sekiguchi et al., 2003) Estimates of the SW, LWS and NET cloud feedbacks are much larger than the other values in the graph.





Figure 7.6 | Change in effective radiative forcing (ERF) from 1750 to 2019 by contributing forcing agents (carbon dioxide, other well-mixed greenhouse gases (WMGHGs), ozone, stratospheric water vapour, surface albedo, contrails and aviation-induced cirrus, aerosols, anthropogenic total, and solar).

Aerosols, climate change and air quality





FAQ 6.2, Figure 1 | Links between actions aiming to limit climate change and actions to improve air quality. Greenhouse gases (GHGs) and aerosols (orange and blue) can affect climate directly. Air pollutants (bottom) can affect human health, ecosystems and climate. All these compounds have common sources and sometimes interact with each other in the atmosphere which makes it impossible to consider them separately (dotted grey arrows).




Figure 6.1 | Sources and processes leading to atmospheric short-lived climate forcer (SLCF) burden and their interactions with the climate system.

FAQ 6.2 | What Are the Links Between Limiting Climate Change and Improving Air Quality?

...]

There are, however, also 'win-lose' actions. For example, wood burning is defined as carbon neutral because a tree accumulates the same amount of carbon dioxide throughout its lifetime as is released when wood from that tree is burned. However, burning wood can also result in significant emissions of air pollutants, including carbon monoxide, nitrogen oxides, volatile organic compounds, and particulate matter, that locally or regionally affect the climate, human health and ecosystems (FAQ 6.2, Figure 1). Alternatively, decreasing the amount of sulphate aerosols produced by power and industrial plants, and from maritime transport, improves air quality but results in a warming influence on the climate, because those sulphate aerosols contribute to cooling the atmosphere by blocking incoming sunlight.

Air quality and climate change represent two sides of the same coin, and addressing both issues together could lead to significant synergies and economic benefits while avoiding policy actions that mitigate one of the two issues but worsen the other.


from MAAP to ASCEND MAAP Service based environment Multi-Mission Algorithm User-centric and Analysis Platform **ASCEND** Configurable **OPEN SOURCE** Advanced Data Access and Secure OPEN COLLABORATION Processing Services for Collaborative EO grouND segment **OPEN SCIENCE** Infrastructure agnostic Distributed **OPEN DATA** Powering the EO SCIENCE community with data, resources, tools and support **Accelerating** EO science **CLOUD COMPUTING** to make advancements **REAL-TIME COLLABORATIVE** Boosting the impact of workspace available to the INTERNATIONAL Biomass community. **EOscience** missions

earthcare

Airborne validation campaigns







More than 100 flights (... and counting)!

International effort involving complex cooperation between more than 60 research organisations, space agencies and aircraft teams

Legacy, Future and Internat'l Collaboration









AOS-Storm/PMM
2029AOS-Sky/HAWCsat
2031-





ACDL/DQ-1, 2022-532nm(HSR)/1064nm/ 1572nm Lidar ACDL/DQ-2, 2025-

CHINA NATIONAL SPACE ADMINISTRATION



eps-Aeolus/-2 2034-355nm HSRL Lidar























